

## Package Leaflet: Information for the User

### **Cutivate 0.05% w/w Cream** fluticasone propionate

**Read all of this leaflet carefully before you start using this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

#### **What is in this leaflet:**

- 1 What Cutivate is and what it is used for
- 2 What you need to know before you use Cutivate
- 3 How to use Cutivate
- 4 Possible side effects
- 5 How to store Cutivate
- 6 Contents of the pack and other information

## **1. What Cutivate is and what it is used for**

Cutivate contains a medicine called fluticasone propionate. It belongs to a group of medicines called steroids. It helps to reduce swelling and irritation.

Cutivate is used to help reduce the redness and itchiness of certain skin problems.

- For adults these skin problems include eczema, psoriasis, dermatitis, insect bites or 'prickly heat'.
- For infants and children it is used for dermatitis that has not responded to milder steroid creams or ointments, following doctors advice.

## **2. What you need to know before you use Cutivate**

#### **Do not use Cutivate:**

- if you are allergic to fluticasone propionate or any of the other ingredients of this medicine (listed in Section 6)
- on an infant under 3 months
- to treat any of the following skin problems, it could make them worse:
  - acne
  - severe flushing of skin on and around your nose (rosacea)
  - spotty red rash around your mouth (perioral dermatitis)
  - itching around your back passage or private parts
  - infected skin (unless the infection is being treated with an anti-infective medicine at the same time)
  - itchy skin which is not inflamed

Do not use if any of the above apply to you. If you are not sure, talk to your doctor or pharmacist before using Cutivate.

#### **Warnings and precautions**

Talk to your doctor or pharmacist before using Cutivate if:

- you have previously had an allergic reaction with another steroid
- you are applying the cream under an airtight dressing, including a child's nappy. These dressings make it easier for the active ingredient to pass through the skin. It is possible to accidentally end up using too much cream.
- you are applying the cream on broken or damaged skin or within skin folds.
- you are applying to a large surface area
- you have psoriasis, your doctor will want to see you more often.
- you are using for a chronic leg ulcer as you may be at increased risk of local reaction or infection
- you are applying near eyes or on eyelids, as cataracts or glaucoma may result if the cream repeatedly enters the eye
- you have an infection of the skin as this will need to be treated
- you are applying to thin skin such as the face, as Cutivate may cause skin thinning.

If you are not sure if the above apply to you, talk to your doctor or pharmacist before using this medicine.

Contact your doctor if you experience blurred vision or other visual disturbances.

### **Children**

Avoid continuous treatment for a long period of time in infants and children under 12 years of age, as their skin may absorb larger amounts.

Whilst using Cutivate, do not smoke or go near naked flames due to the risk of severe burns. Fabric (clothing, bedding, dressing etc) that has been in contact with this product burns more easily and is a serious fire hazard. Washing clothing and bedding may reduce product build-up but not totally remove it.

If there is a worsening of your condition during use consult your prescriber – you may be experiencing an allergic reaction, have an infection or your condition requires a different treatment.

If you experience a recurrence of your condition shortly after stopping treatment, within 2 weeks, do not restart using the cream/ointment without consulting your prescriber unless your prescriber has previously advised you to so. If your condition has resolved and on recurrence the redness extends beyond the initial treatment area and you experience a burning sensation, please seek medical advice before restarting treatment.

### **Other medicines and Cutivate**

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicine, especially if you are taking ritonavir and itraconazole medications.

### **Pregnancy, breast-feeding and fertility**

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

**Cutivate Cream contains imidurea, cetostearyl alcohol and propylene glycol**

Cutivate cream contains an ingredient called imidurea. Your body can break down imidurea in to small amounts of a chemical called formaldehyde. This chemical may cause a skin reaction, including redness and itchiness.

Cutivate cream also contains cetostearyl alcohol which may cause local skin reactions (e.g. local dermatitis) and 100 mg propylene glycol in each gram which may cause local skin irritation.

### 3. How to use Cutivate

Always use this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

#### Using this medicine

- Apply a thin layer of Cutivate up to 2 times a day. This may be reduced as your skin begins to get better.
- This cream is for use on your skin only.
- Do not use on large areas of the body for a long time (such as every day for many weeks or months) - unless your doctor tells you to.
- The germs that cause infections like warm and moist conditions under dressings so always clean the skin before a fresh dressing is put on.
- If you are applying the cream on someone else make sure you wash your hands after use or wear disposable plastic gloves.
- If your skin problem does not improve, talk to your doctor.

#### Guidance on how to apply the cream

- 1 Wash your hands.
- 2 Apply a thin layer to the affected area(s) and gently rub into the skin until it has all disappeared. You can measure how much Cutivate to use with your fingertip. For children you will need to use less cream but still use an adult finger to measure out the fingertip unit. This picture shows one fingertip unit.

#### [Pictogram of 1 fingertip unit]

- 3 Unless you are meant to apply the cream to your hands as a part of the treatment, wash them again after using the cream.

#### For an adult

#### [Pictogram of an adult with fingertip units to be used as per text below with arrows pointing to relevant body part]

Face and neck:

**2 ½ fingertip units.**

Back: **7 fingertip units.**

Front: **7 fingertip units.**

One arm (not including the hand):

**3 fingertip units.**

Both sides of one hand:

**1 fingertip unit.**

One leg (not including the foot):  
**6 fingertip units**

One foot:  
**2 fingertip units.**

Do not worry if you find you need a little more or less than this. It is only a rough guide.

#### For a child

Number of fingertip units needed					
Child's age	Face and neck	Arm and hand	Leg and foot	Front	Back including buttocks
3-6 months	1	1	1 ½	1	1 ½
1-2 years	1 ½	1 ½	2	2	3
3-5 years	1 ½	2	3	3	3 ½
6-10 years	2	2 ½	4 ½	3 ½	5

- Do not use it on infants under 3 months of age.
- A doctor will provide expert advice on treating your child's skin problem.
- If the child's skin problem does not improve in 7 to 14 days, talk to your doctor.
- It should not be used every day for more than 4 weeks.

#### **If you apply Cutivate to your face**

You should only apply the cream to your face if your doctor tells you to. The cream should not be used for too long as the skin on your face thins easily. **Do not let the cream get into your eyes.**

#### **If you use more Cutivate than you should**

If, by mistake on a few occasions you use more than you should, do not worry. If you apply a lot or if a lot is accidentally swallowed, it could make you ill. Talk to your doctor or go to hospital as soon as possible.

#### **If you forget to use Cutivate**

If you forget to apply your cream, apply it as soon as you remember. If it is close to the time you are next meant to apply it, wait until this time.

#### **If you stop using Cutivate**

If you use Cutivate regularly make sure you talk to your doctor before you stop using it.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

#### 4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

**Stop using Cutivate and tell your doctor as soon as possible if:**

- you find that your skin problem gets worse or becomes swollen during treatment. You may be allergic to the cream, have an infection or need other treatment.

**Other side effects you may notice when using Cutivate include:**

**Common** (may affect up to 1 in 10 people)

- Irritation or itching.

**Uncommon** (may affect up to 1 in 100 people)

- A feeling of burning where the cream is applied.

**Very rare** (may affect up to 1 in 10,000 people)

- If you have psoriasis you may get raised bumps with pus under the skin. This can happen during or after the treatment and is known as pustular psoriasis.
- Worsening of condition
- Redness
- Rash or hives
- Stretch marks may develop
- Blood vessels under the surface of your skin may become more noticeable
- Changes to the colour of your skin
- Increased body hair
- Skin thinning
- Skin wrinkling
- Skin dryness
- Weight gain and rounding of the face
- Obesity

In children, also look out for the following symptoms:

- Delayed weight gain
- Slow growth

Very rare side effects that may show up in blood tests or when your doctor gives you a medical examination:

- A decrease in the level of the hormone cortisol in your blood
- Increased levels of sugar in your blood or sugar in the urine
- High blood pressure
- Cloudy lens in the eye (cataract)
- Increased pressure in the eye (glaucoma)

- Weakening of the bones through gradual loss of mineral (osteoporosis); additional tests may be needed after your medical examination to confirm if you have this condition

**Not Known** (frequency cannot be estimated from the available data)

- Blurred vision
- Steroid withdrawal reaction: If used continuously for prolonged periods a withdrawal reaction may occur on stopping treatment with some or all of the following features: redness of the skin which can extend beyond the initial area treated, a burning or stinging sensation, intense itching, peeling of the skin, oozing open sores

### **Reporting of side effects**

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App Store.

By reporting side effects you can help provide more information on the safety of this medicine.

## **5. How to store Cutivate**

- Keep this medicine out of the sight and reach of children.
- Do not use this medicine after the expiry date which is stated on the tube or carton after EXP. The expiry date refers to the last day of that month.
- Do not store above 30°C.
- Do not throw away medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

## **6. Contents of the pack and other information**

### **What Cutivate contains**

- The active ingredient is fluticasone propionate. Each 1 g contains 0.5 mg of fluticasone propionate (0.05% w/w).
- The other ingredients are liquid paraffin (see section 2), cetostearyl alcohol (see section 2), isopropyl myristate, cetomacrogol 1000, propylene glycol (see section 2), imidurea (see section 2), sodium phosphate, citric acid monohydrate and purified water.

### **What Cutivate looks like and contents of the pack**

Within each carton is a tube with a plastic screw cap, which contains either 15, 30 or 50 g of cream. Not all pack sizes may be marketed.

### **Marketing Authorisation Holder and Manufacturer**

Product licence held by Glaxo Wellcome UK Ltd, GSK Medicines Research Centre, Gunnels Wood Road, Stevenage, Hertfordshire, SG1 2NY, UK  
 Manufactured by Glaxo Operations UK Ltd (trading as Glaxo Wellcome Operations), Harmire Road, Barnard Castle, Durham, DL12 8DT

**More Information**

If you have any questions or are not sure about anything, ask your doctor or pharmacist who will advise you.

You may be able to find out more from books in public libraries.

**Other formats**

To listen to or request a copy of this leaflet in Braille, large print or audio please call, free of charge:

0800 198 5000 (UK Only)

Please be ready to give the following information:

**Product name**      **Cutivate 0.05% w/w Cream**

Reference number 10949/0013

This is a service provided by the Royal National Institute of Blind People.

**This leaflet was last revised in August 2023**

Trade marks are owned by or licensed to the GSK group of companies

© 2023 GSK group of companies or its licensor