

## Package leaflet: Information for the user

### Micafungin 50 mg and 100 mg Powder for Concentrate for Solution for Infusion

**Read all of this leaflet carefully before you start using this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- If you get any side effects, talk to your doctor, or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

#### What is in this leaflet

1. **What Micafungin is and what it is used for**
2. **What you need to know before you use Micafungin**
3. **How to use Micafungin**
4. **Possible side effects**
5. **How to store Micafungin**
6. **Contents of the pack and other information**

#### 1. What Micafungin is and what it is used for

Micafungin contains the active substance micafungin. Micafungin is called an antifungal medicine because it is used to treat infections caused by fungal cells.

Micafungin is used to treat fungal infections caused by fungal or yeast cells called Candida.

Micafungin is effective in treating systemic infections (those that have penetrated within the body). It interferes with the production of a part of the fungal cell wall. An intact cell wall is necessary for the fungus to continue living and growing. Micafungin causes defects in the fungal cell wall, making the fungus unable to live and grow.

Your doctor has prescribed Micafungin for you in the following circumstances when there are no other suitable antifungal treatments available (see section 2):

- To treat adults, adolescents and children including neonates who have a serious fungal infection called invasive candidiasis (infection that has penetrated the body).
- To treat adults and adolescents  $\geq 16$  years of age who have a fungal infection in the gullet (oesophagus) where treatment into a vein (intravenous) is appropriate.
- To prevent infection with Candida in patients who are having a bone-marrow transplant or who are expected to have a neutropenia (low levels of neutrophils, a type of white blood cell) for 10 days or more.

#### 2. What you need to know before you use Micafungin

##### Do not use Micafungin:

- if you are allergic to micafungin, other echinocandins (Ecalta or Cancidas) or any of the other ingredients of this medicine (listed in section 6).

#### Warnings and precautions

In rats, long-term treatment with micafungin led to liver damage and subsequent liver tumours. The potential risk of developing liver tumours in humans is not known, and your doctor will assess the benefits and risks of Micafungin treatment before starting your medicine. Please tell your doctor if you have severe liver problems (e.g. liver failure or hepatitis) or have had abnormal liver function tests.

During treatment your liver functions will be monitored more closely.

Talk to your doctor or, pharmacist before using Micafungin

- if you are allergic to any medicine
- if you have haemolytic anaemia (anaemia due to breakdown of red blood cells) or haemolysis (breakdown of red blood cells)
- if you have kidney problems (e.g. kidney failure and abnormal kidney function test). If this happens, your doctor may decide to monitor your kidney function more closely.

Micafungin may also cause severe inflammation/eruption of the skin and mucous membranes (Stevens-Johnson syndrome, toxic epidermal necrolysis).

### **Other medicines and Micafungin**

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

It is especially important to inform your doctor if you are using amphotericin B desoxycholate or itraconazole (antifungal antibiotics), sirolimus (an immunosuppressant) or nifedipine (calcium channel blocker used to treat high blood pressure). Your doctor may decide to adjust the dose of these medicines.

### **Micafungin with food and drink**

As Micafungin is given intravenously (into a vein), no restrictions on food or drink are required.

### **Pregnancy and breast-feeding**

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

Micafungin should not be used during pregnancy unless clearly necessary. If you use Micafungin you should not breast-feed.

### **Driving and using machines**

Micafungin is unlikely to have an effect on driving or using machines. However some people may feel dizzy when taking this medicine and if this happens to you, do not drive or use any tools or machines. Please inform your doctor if you experience any effects that may cause you to have problems with driving or using other machinery.

### **Micafungin contains sodium**

Micafungin contains less than 1 mmol sodium (23 mg) per vial, that is to say essentially 'sodium-free'.

## **3. How to use Micafungin**

Micafungin must be prepared and given to you by a doctor or another healthcare professional.

Micafungin should be administered once daily by slow intravenous (into a vein) infusion. Your doctor will determine how much Micafungin you will receive each day.

### **Use in adults, adolescents $\geq$ 16 years of age and elderly**

- The usual dose to treat an invasive Candida infection is 100 mg per day for patients weighing more than 40 kg and 2 mg/kg per day for patients weighing 40 kg or less.
- The dose to treat a Candida infection of the oesophagus is 150 mg for patients weighing more than 40 kg and 3 mg/kg per day for patients weighing 40 kg or less.
- The usual dose to prevent invasive Candida infections is 50 mg per day for patients weighing more than 40 kg and 1 mg/kg per day for patients weighing 40 kg or less.

### **Use in children > 4 months of age and adolescents < 16 years of age**

- The usual dose to treat an invasive Candida infection is 100 mg per day for patients weighing more than 40 kg and 2 mg/kg per day for patients weighing 40 kg or less.

- The usual dose to prevent invasive Candida infections is 50 mg per day for patients weighing more than 40 kg and 1 mg/kg per day for patients weighing 40 kg or less.

#### **Use in children and newborns < 4 months of age**

- The usual dose to treat an invasive Candida infection is 4-10 mg/kg per day.
- The usual dose to prevent invasive Candida infections is 2 mg/kg per day.

#### **If you receive more Micafungin than you should**

Your doctor monitors your response and condition to determine what dose of Micafungin is needed. However, if you are concerned that you may have been given too much Micafungin, speak to your doctor or another healthcare professional immediately.

#### **If you miss a dose of Micafungin**

Your doctor monitors your response and condition to determine what Micafungin treatment is needed. However, if you are concerned that you may have missed a dose, speak to your doctor or another healthcare professional immediately.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

## **4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

If you experience an allergic attack, or a severe skin reaction (e.g. blistering and peeling of the skin), you must inform your doctor or nurse immediately.

Micafungin may cause the following other side effects:

#### **Common (may affect up to 1 in 10 people)**

- abnormal blood tests (decreased white blood cells [leucopenia; neutropenia]); decreased red blood cells (anaemia)
- decreased potassium in the blood (hypokalaemia); decreased magnesium in the blood (hypomagnesaemia); decreased calcium in the blood (hypocalcaemia)
- headache
- inflammation of the vein wall (at injection-site)
- nausea (feeling sick); vomiting (being sick); diarrhoea; abdominal pain
- abnormal liver function tests (increased alkaline phosphatase; increased aspartate aminotransferase, increased alanine aminotransferase)
- increased bile pigment in the blood (hyperbilirubinaemia)
- rash
- fever
- rigors (shivering).

#### **Uncommon (may affect up to 1 in 100 people)**

- abnormal blood tests (decreased blood cells [pancytopenia]); decreased blood platelets (thrombocytopenia); increases in a certain type of white blood cells called eosinophils; decreased albumin in the blood (hypoalbuminaemia)
- hypersensitivity
- increased sweating
- decreased sodium in the blood (hyponatraemia); increased potassium in the blood (hyperkalaemia); decreased phosphates in the blood (hypophosphataemia); anorexia (eating disorder)
- insomnia (difficulty in sleeping); anxiety; confusion
- feeling lethargic (somnolence); trembling; dizziness; disturbed taste
- increased heart rate; stronger heartbeat; irregular heartbeat

- high or low blood pressure; skin flushing
- shortness of breath
- indigestion; constipation
- liver failure; increased liver enzymes (gamma-glutamyltransferase); jaundice (yellowing of the skin or whites of the eyes caused by liver or blood problems); reduced bile reaching the intestine (cholestasis); enlarged liver; liver inflammation
- itchy rash (urticaria); itching; skin flushing (erythema)
- abnormal kidney function tests (increased blood creatinine; increased urea in the blood); aggravated kidney failure
- increase in an enzyme called lactate dehydrogenase
- clotting in vein at injection-site; inflammation at injection-site; pain at injection-site; collection of fluid in your body.

**Rare (may affect up to 1 in 1,000 people)**

- anaemia due to breakdown of red blood cells (haemolytic anaemia), breakdown of red blood cells (haemolysis).

**Not known (frequency cannot be estimated from the available data)**

- disorder of blood clotting system
- (allergic) shock
- damage to liver cells including death
- kidney problems; acute kidney failure.

**Additional side effects in children and adolescents**

The following reactions have been reported more often in paediatric patients than in adult patients:

**Common (may affect up to 1 in 10 people)**

- decreased blood platelets (thrombocytopenia)
- increased heart rate (tachycardia)
- high or low blood pressure
- increased bile pigment in the blood (hyperbilirubinaemia); enlarged liver
- acute kidney failure; increased urea in the blood.

**Reporting of side effects**

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme Website: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

**5. How to store Micafungin**

**Keep this medicine out of the sight and reach of children.**

Do not use this medicine after the expiry date which is stated on the vial and on the carton. The expiry date refers to the last day of that month.

The unopened vial does not require any special storage conditions.

**Reconstituted concentrate in vial**

Chemical and physical in-use stability has been demonstrated for up to 48 hours at 25°C when reconstituted with sodium chloride 9 mg/ml (0.9%) solution for infusion or glucose 50 mg/ml (5%) solution for infusion.

**Diluted infusion solution**

Chemical and physical in-use stability has been demonstrated for 96 hours at 25°C when protected from light when diluted with sodium chloride 9 mg/ml (0.9%) solution for infusion or glucose 50 mg/ml (5%) solution for infusion.

From a microbiological point of view, the reconstituted and diluted solutions should be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would normally not be longer than 24 hours at 2 to 8°C, unless the reconstitution and dilution have taken place in controlled and validated aseptic conditions.

Only a trained healthcare professional who has read the complete directions properly can prepare this medicine for use.

Do not use the diluted infusion solution if it is cloudy or precipitated.

In order to protect the infusion bottle / bag containing the diluted infusion solution from light it should be inserted into a closable opaque bag.

The vial is for single use only. Therefore, please discard unused reconstituted concentrate immediately.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

## **6. Contents of the pack and other information**

### **What Micafungin contains**

- The active substance is micafungin (as sodium).
- 1 vial contains micafungin sodium equivalent to 50 mg or 100 mg micafungin.
- The other ingredients are sucrose, citric acid and sodium hydroxide.

### **What Micafungin looks like and contents of the pack**

Micafungin 50 mg or 100 mg powder for concentrate for solution for infusion is a white to off-white powder.

Micafungin is supplied in a box containing 1 vial.

### **Marketing Authorisation Holder**

TEVA UK Limited, Ridings Point, Whistler Drive, Castleford, WF10 5HX, United Kingdom

### **Manufacturer**

S.C. SINDAN-PHARMA S.R.L., 11, Ion Mihalache Ave., the 1st district, Bucharest, 011171, Romania

OR\*

Merckle GmbH, Graf-Arco-Str. 3, 89079 Ulm, Germany

**This leaflet was last revised in March 2024**

PL 00289/2322 & PL 00289/2323

*\*Only the actual site of batch release will appear on the printed version of the leaflet.*

-----

--

The following information is intended for medical or healthcare professionals only:

Micafungin must not be mixed or co-infused with other medicinal products except those mentioned below. Using aseptic techniques at room temperature, Micafungin is reconstituted and diluted as follows:

1. The plastic cap must be removed from the vial and the stopper disinfected with alcohol.
2. Five ml of sodium chloride 9 mg/ml (0.9%) solution for infusion or glucose 50 mg/ml (5%) solution for infusion (taken from a 100 ml bottle/bag) should be aseptically and slowly injected into each vial along the side of the inner wall. Although the concentrate will foam, every effort should be made to minimise the amount of foam generated. A sufficient number of vials of Micafungin must be reconstituted to obtain the required dose in mg (see table below).
3. The vial should be rotated gently. **DO NOT SHAKE**. The powder will dissolve completely. The concentrate should be used immediately. The vial is for single use only. Therefore, unused reconstituted concentrate must be discarded immediately.
4. All of the reconstituted concentrate should be withdrawn from each vial and returned into the infusion bottle/bag from which it was originally taken. The diluted infusion solution should be used immediately. Chemical and physical in-use stability have been demonstrated for 96 hours at 25°C when protected from light and diluted as described above.
5. The infusion bottle/bag should be gently inverted to disperse the diluted solution but **NOT** agitated in order to avoid foaming. The solution must not be used if it is cloudy or has precipitated.
6. The infusion bottle/bag containing the diluted infusion solution should be inserted into a closable opaque bag for protection from light.

Preparation of the solution for infusion

<b>Dose (mg)</b>	<b>Micafungin vial to be used (mg/vial)</b>	<b>Volume of sodium chloride (0.9%) or glucose (5%) to be added per vial</b>	<b>Volume (concentration) of reconstituted powder</b>	<b>Standard infusion (added up to 100 ml) final concentration</b>
50	1 x 50	5 ml	approx. 5 ml (10 mg/ml)	0.5 mg/ml
100	1 x 100	5 ml	approx. 5 ml (20 mg/ml)	1.0 mg/ml
150	1 x 100 + 1 x 50	5 ml	approx. 10 ml	1.5 mg/ml
200	2 x 100	5 ml	approx. 10 ml	2.0 mg/ml

After reconstitution and dilution, the solution should be administered by intravenous infusion over approximately 1 hour.