

<b>BOZZA N°</b>	01 06.10.22	02 10.10.22	03 05.06.23	04 29.06.23	05 14.08.23					
-----------------	----------------	----------------	----------------	----------------	----------------	--	--	--	--	--

Fenofibrate\_267 mg hard capsules Leaflet Fto: 175x 290 mm

Font e size utilizzate:

helvetica LT Std condensed: c. 8,3 / 9,52

helvetica LT Std Bold condensed: c. 8,3 / 9,52

175 mm

**Package leaflet: Information for the patient**  
**Fenofibrate 267mg capsules**  
Fenofibrate

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- If any of the side effects get serious, or if you notice any side effects not listed in this leaflet, please tell your doctor, or pharmacist or nurse.

**What is in this leaflet:**

1. What Fenofibrate is and what it is used for
2. What you need to know before you take Fenofibrate
3. How to take Fenofibrate
4. Possible side effects
5. How to store Fenofibrate
6. Contents of the pack and other information

**1. What Fenofibrate is and what it is used for**  
Fenofibrate belongs to a group of medicines, commonly known as 'fibrates'. These medicines are used to lower the level of fats (lipids) in the blood. For example the fats known as 'triglycerides'.

Fenofibrate is used, alongside a low fat diet and other non-medical treatments such as exercise and weight loss, to lower levels of fats in the blood.

Fenofibrate can be used in addition to other medicines (called 'statins') in some circumstances when levels of fats in the blood are not controlled with a statin alone.

Fenofibrate can often also increase the amount of a 'good' type of cholesterol, called HDL or high density lipoprotein cholesterol.

It is always essential to continue a low-fat diet during treatment with Fenofibrate.

**2. What you need to know before you take Fenofibrate**  
Do not take Fenofibrate if:

- You are allergic to fenofibrate or any of the other ingredients of this medicine as it may cause allergic reactions (possibly delayed) (listed in Section 6: contents of the pack and other information)
- If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.
- While taking other medicines, you have had an allergic reaction or skin damage from sunlight or UV light (these medicines include other fibrates and an anti-inflammatory medicine called 'ketoprofen')
- You have severe liver, kidney or gallbladder problems
- You have pancreatitis (an inflamed pancreas which causes abdominal pain), which is not caused by high levels of fat in the blood

Do not take Fenofibrate if any of the above apply to you. If you are not sure, talk to your doctor or pharmacist before taking Fenofibrate.

**Warnings and Precautions**  
Talk to your doctor or pharmacist or nurse before taking Fenofibrate if:

- You have any liver or kidney problems
- You may have an inflamed liver (hepatitis) - signs include yellowing of the skin and the whites of the eyes Jaundice) and an increase in liver enzymes (shown in blood tests)
- You have an under-active thyroid gland (hypo-thyroidism)
- You have diabetes, especially Type 2 diabetes, that is not well controlled
- You have problems with certain proteins in your blood
- You have an alcohol problem
- You are taking other medicines
- You or your family have had muscle problems
- You are over 70 years of age

(Some of the above conditions can lead to high levels of lipids in your blood and need to be corrected before you start therapy with fenofibrate).

If any of the above apply to you (or you are not sure), talk to your doctor or pharmacist before taking Fenofibrate.

Your doctor might want to test your blood or urine to check if Fenofibrate is working properly and if your kidneys, muscles and liver are working properly.

**Effects on muscles**  
Stop taking Fenofibrate and see a doctor straight away if you get unexplained cramps or painful, tender or weak muscles while taking this medicine.

- This is because this medicine may cause muscle problems, which may be serious.
- These problems are rare but include muscle inflammation and breakdown. This can cause kidney damage or even death.

Your doctor may do a blood test to check your muscles before and after starting treatment.

The risk of muscle breakdown is higher in some patients. Tell your doctor if:

- You are over 70 years old
- You have kidney problems
- You have thyroid problems
- You or a close family member has a muscle problem which runs in the family
- You drink large amounts of alcohol
- You are taking medicines called statins to lower cholesterol - such as simvastatin, atorvastatin, pravastatin, rosuvastatin or fluvastatin
- You have ever had muscle problems during treatment with statins or fibrates - such as fenofibrate, bezafibrate or gemfibrozil

If any of the above apply to you (or you are not sure), talk to your doctor before taking Fenofibrate.

Fenofibrate Capsules contain sucrose  
This medicine contains sucrose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

Fenofibrate Capsules contain methyl p-hydroxybenzoate and propyl phydroxybenzoate. This medicine contains methyl p-hydroxybenzoate and propyl p-hydroxybenzoate, which may cause allergic reactions (possibly delayed).

Fenofibrate Capsules contains less than 1 mmol (23 mg) sodium per capsule, that is to say essentially sodium-free.

**Other medicines and Fenofibrate**  
Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription.  
In particular tell your doctor or pharmacist if you are taking any of the following medicines:

- Anti-coagulants to thin your blood (such as warfarin)
- Other medicines to control fat levels in the blood (such as 'statins' or 'fibrates'). Taking a 'statin' at the same time as Fenofibrate may increase the risk of muscle problems
- A particular class of medicines to treat diabetes (such as rosiglitazone or pioglitazone)
- Cyclosporin - used to suppress your immune system

If any of the above apply to you (or you are not sure), talk to your doctor or pharmacist before taking Fenofibrate.

**Fenofibrate with food, drink and alcohol**  
It is important to take the capsule with food - it will not work as well if your stomach is empty.

**Pregnancy, breast-feeding and fertility**

- Do not take Fenofibrate and tell your doctor if you are pregnant, think you might be pregnant or are planning to have a baby
- Do not take Fenofibrate if you are breast-feeding or planning to breast-feed your baby

Ask your doctor or pharmacist for advice before taking any medicine.

**Driving and using machines**  
This medicine will not affect you being able to drive or use tools or machines.

**3. How to take Fenofibrate**  
Always take this medicine exactly as your doctor or pharmacist has told you. Also, please read the label on the packet. You should check with your doctor or pharmacist if you are not sure.

Your doctor will determine the appropriate strength for you, depending on your condition, your current treatment and your personal risk status.

290 mm

FRONT

290 mm

BACK

175 mm

**Taking this medicine**

- Swallow the capsule whole with a glass of water
- Do not open or chew the capsule
- Take the capsule with food - it will not work as well if your stomach is empty

**How much to take**  
The recommended dose for adults is one capsule of Fenofibrate 200 mg a day, taken at mealtimes. However, your doctor may want you to take one capsule of Fenofibrate 267mg a day (a higher dose).

**Use in children and adolescents**  
The use of Fenofibrate 267mg is not recommended in children under the age of 18.

**People with kidney problems**  
If you have kidney problems, your doctor may tell you to take a lower dose. Ask your doctor or pharmacist about this.

**If you take more Fenofibrate than you should**  
If you take more Fenofibrate than you should or if someone else has taken your medicine, contact your nearest hospital casualty department or tell your doctor immediately.

**If you forget to take Fenofibrate**

- If you forget a dose, take the next dose with your next meal
- Then take your next capsule at the normal time
- Do not take a double dose to make up for a forgotten dose.

If you are worried about this talk to your doctor.

If you stop taking Fenofibrate  
Do not stop taking Fenofibrate unless your doctor tells you to, or the capsules make you feel unwell. This is because abnormal levels of fats in the blood need treating for a long period of time.

Remember that as well as taking Fenofibrate it is also important that you:

- Have a low fat diet
- Take regular exercise

If you have any further questions on the use of this product, ask your doctor or pharmacist or nurse.

**4. Possible side effects**  
Like all medicines, this medicine can cause side effects, although not everybody gets them.

**Stop taking Fenofibrate and see a doctor straight away, if you notice any of the following serious side effects - you may need urgent medical treatment:**

- Allergic reaction - the signs may include swelling of the face, lips, tongue or throat, which may cause difficulty in breathing
- Cramps or painful, tender or weak muscles - these may be signs of muscle inflammation or breakdown, which can cause kidney damage or even death
- Stomach pain - this may be a sign that your pancreas is inflamed (pancreatitis)
- Chest pain and feeling breathless - these may be signs of a blood clot in the lung (pulmonary embolism)
- Pain, redness or swelling in the legs - these may be signs of a blood clot in the leg (deep vein thrombosis)
- Yellowing of the skin and whites of the eyes Jaundice), or an increase in liver enzymes - these may be signs of an inflamed liver (hepatitis)

Stop taking Fenofibrate and see a doctor straight away, if you notice any of the side effects above.

**Other side effects include:**  
**Common** (affects less than 1 in 10 people):

- Diarrhoea
- Stomach pain
- Wind (flatulence)
- Feeling sick (nausea)
- Being sick (vomiting)
- Raised levels of liver enzymes in the blood - shown in tests
- Increase in homocysteine (too much of this amino acid in the blood has been associated to a higher risk of coronary heart disease, stroke and peripheral vascular disease, although a causal link has not been established)

**Uncommon** (affects less than 1 in 100 people):

- Headache
- Gallstones

- Reduced sex drive
- Rash, itching or red patches on the skin
- Increase in 'creatinine' produced by the kidneys - shown in tests
- Pancreatitis (inflammation of the pancreas leading to abdominal pain)
- Thromboembolism: pulmonary embolism (blood clot in the lung causing chest pain and breathlessness), deep vein thrombosis (blood clot in the leg causing pain, redness or swelling)
- Muscle pain, muscle inflammation, muscle cramps and weakness

**Rare (affects less than 1 in 1,000 people):**

- Hair loss
- Increase in 'urea' produced by the kidneys - shown in tests
- Increased sensitivity of your skin to sunlight, sun lamps and sunbeds
- Drop in haemoglobin (that carries oxygen in the blood) and white blood cells - shown in tests.
- Hepatitis (inflammation of the liver), symptoms of which may be mild jaundice (yellowing of the skin and whites of the eyes), stomach pain and itching
- Hypersensitivity (allergic reaction)

**Side effect where the chance of it happening are not known**

- Severe form of skin rash with reddening, peeling and swelling of the skin that resembles severe burns
- Long-term lung problems
- Muscle breakdown
- Complications of gallbladder stones
- Jaundice
- Feeling dizzy (vertigo)
- Feeling exhausted (fatigue)

If you get any unusual breathing discomfort, tell your doctor straight away.  
If you get any side effects, talk to your doctor, or pharmacist or nurse. This includes any side effects not listed in this leaflet.

**Reporting of side effects**  
If you get any side effects, talk to your doctor or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

**5. How to store Fenofibrate**  
Keep this medicine out of the sight and reach of children.  
This medicinal product does not require any special storage conditions. Do not use this medicine after the expiry date which is stated on the carton and the blister after EXP. The expiry date refers to the last day of that month.  
Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

**6. Contents of the pack and other information**  
**What Fenofibrate capsules contains**

- The active substance is fenofibrate. Each Fenofibrate 267mg capsule contains 267 milligrams (mg) of micronised fenofibrate. This medicine contains 0.011 mg sodium benzoate in each capsule.
- The other ingredients are: sugar spheres, hypromellose, sodium lauril sulfate, talc, polydimethylsiloxane, octylphenoxy polyethoxy ethanol, polyethylene glycol sorbitan monolaurate, propylene glycol (E 1520), sodium benzoate (E 211), methyl p-hydroxybenzoate (E 218), propyl p-hydroxybenzoate (E 216), simethicone, cetyl stearyl alcohol and ethoxylate.

The capsule is made of gelatin, titanium dioxide (E171), yellow iron oxide (E172).

**What Fenofibrate capsules looks like and contents of the pack**  
Fenofibrate 267mg capsules are supplied to you as yellow/transparent hard capsules. Fenofibrate 267mg capsules are provided in blister packs of 28 or 30 capsules. Not all pack sizes may be marketed.

**Marketing Authorisation Holder and Manufacturer:**

Rivopharm UK Ltd  
100 Bishopsgate  
London, EC2N 4AG, United Kingdom

**This leaflet was last revised in Aug 2023.**