

Package leaflet: Information for the user

Anagrelide 0.5 mg hard capsules

Anagrelide hydrochloride monohydrate

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet:

1. What is Anagrelide and what it is used for
2. What you need to know before you take Anagrelide
3. How to take Anagrelide
4. Possible side effects
5. How to store Anagrelide
6. Contents of the pack and other information.

1. What is Anagrelide and what it is used for

Anagrelide contains the active substance, anagrelide. Anagrelide is a medicine which interferes with the development of platelets. It reduces the number of platelets produced by the bone marrow, which results in a decrease in the platelet count in the blood towards a more normal level. For this reason it is used to treat patients with essential thrombocythaemia.

Essential thrombocythaemia is a condition which occurs when the bone marrow produces too many of the blood cells known as platelets. Large numbers of platelets in the blood can cause serious problems with blood circulation and clotting.

2. What you need to know before you take Anagrelide

Do not take Anagrelide

- If you are allergic to anagrelide or any of the other ingredients of this medicine (listed in section 6). An allergic reaction may be recognized as a rash, itching, swollen face or lips, or shortness of breath;
- If you have moderate or severe liver problems;
- If you have moderate or severe kidney problems.

Warnings and precautions

Talk to your doctor **before taking Anagrelide**

- If you have or think you might have a problem with your heart;
- If you were born with or have family history of prolonged QT interval (seen on ECG, electrical recording of the heart), or you are taking other medicines that result in abnormal ECG changes or if you have low levels of electrolytes e.g. potassium, magnesium or calcium (see section "Other medicines and Anagrelide")
- If you have any problems with your liver or kidneys.

In combination with acetylsalicylic acid (a substance present in many medicines used to relieve pain and lower fever, as well as to prevent blood clotting, also known as aspirin), there is an increased risk of major haemorrhages (bleeding) (see section "Other medicines and Anagrelide").

While taking Anagrelide, you should take the exact dose prescribed by your doctor. Do not stop taking the medicine without first talking to your doctor. Do not abruptly stop taking this medicine without consulting your doctor. Abrupt withdrawal of medicine may lead to increased risk of stroke.

Signs and symptoms of stroke may include sudden numbness or weakness in the face, arm, or leg, especially on one side of the body, sudden confusion, trouble speaking, or difficulty understanding speech, sudden trouble seeing in one or both eyes, sudden trouble walking, dizziness, loss of balance, or lack of coordination and sudden severe headache with no known cause. Please seek immediate medical help.

Children and adolescents

There is limited information on the use of Anagrelide in children and adolescents and therefore this medicine should be used with caution.

Other medicines and Anagrelide

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Tell your doctor if you are taking any of the following medicines:

- Medicines that can alter your heart rhythm e.g. sotalol, amiodarone;
- Fluvoxamine, used to treat depression;

- Certain types of antibiotic, such as enoxacin, used to treat infections;
- Theophylline, used to treat severe asthma and breathing problems;
- Medicines used to treat heart disorders, for example, milrinone, enoximone, amrinone, olprinone and cilostazol;
- Acetylsalicylic acid (a substance present in many medicines used to relieve pain and lower fever, as well as to prevent blood clotting, also known as aspirin);
- Other medicines used to treat conditions affecting the platelets in your blood, e.g. clopidogrel;
- Omeprazole, used to reduce the amount of acid produced in the stomach;
- Oral contraceptives: If you experience bad diarrhoea whilst taking this medicine, it may reduce how well the oral contraceptive works and use of an extra method of contraception is recommended (e.g. condom). See the instructions in the patient leaflet of the contraceptive pill you are taking.

Anagrelide or these medicines may not work properly if taken together.

If you are not sure, speak to your doctor or pharmacist for advice.

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. Anagrelide should not be taken by pregnant women. Women who are at risk of becoming pregnant should make sure that they are using effective contraception when taking Anagrelide. Speak to your doctor if you need advice with contraception.

Anagrelide should not be taken while breast-feeding. You must stop breast-feeding if you are taking Anagrelide.

Driving and using machines

Dizziness has been reported by some patients taking Anagrelide. Do not drive or use machines if you feel dizzy.

Anagrelide contains lactose

If you have been told that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

Anagrelide contains sodium

This medicine contains less than 1 mmol sodium (23 mg) per capsule, that is to say essentially 'sodium-free'.

3. How to take Anagrelide

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

The amount of Anagrelide that people take can be different, and this depends on your condition. Your doctor will prescribe the best dose for you.

The usual starting dose of Anagrelide is 1 mg. You take this dose as one capsule of 0.5 mg twice a day, for at least a week. After this time, your doctor may either increase or decrease the number of capsules that you take to find the dose best suited to you and which treats your condition most effectively.

Your capsules should be swallowed whole with a glass of water. Do not crush the capsules or dilute the contents in a liquid. You can take the capsules with food or after a meal or on an empty stomach. It is best to take the capsule(s) at the same time every day.

Do not take more or less capsules than your doctor has recommended. Do not stop taking the medicine without first talking to your doctor. You should not suddenly stop taking this medicine on your own.

Your doctor will ask you to have blood tests at regular intervals to check that your medicine is working effectively and that your liver and kidneys are working well.

If you take more Anagrelide than you should

If you take more Anagrelide than you should or if someone else has taken your medicine, tell a doctor or pharmacist immediately. Show them the pack of Anagrelide.

If you forget to take Anagrelide

Take your capsules as soon as you remember. Take your next dose at the usual time. Do not take a double dose to make up for a forgotten dose.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them. If you are worried, speak to your doctor.

Serious side effects:

Uncommon: Heart failure (signs include shortness of breath, chest pain, swelling of the legs due to fluid build-up), severe problem with the rate or rhythm of the heart beat (ventricular tachycardia, supraventricular tachycardia or atrial fibrillation), inflammation of the pancreas which causes severe abdominal and back pain (pancreatitis),

vomiting blood or passing bloody or black stools, severe reduction in blood cells which can cause weakness, bruising, bleeding or infections (pancytopenia). pulmonary hypertension (signs include shortness of breath, swelling in legs or ankles, and lips and skin can turn bluish colour). Rare: Kidney failure (when you pass little or no urine), heart attack.

If you notice any of these side effects, contact your doctor immediately.

Very common side effects: may affect more than 1 in 10 people
Headache.

Common side effects: may affect up to 1 in 10 people

Dizziness, tiredness, rapid heartbeat, irregular or strong heartbeat (palpitations), feeling sick (nausea), diarrhoea, stomach pain, wind, being sick (vomiting), reduction in red blood cell count (anaemia), fluid retention or rash.

Uncommon side effects: may affect up to 1 in 100 people

A feeling of weakness or feeling unwell, high blood pressure, irregular heartbeat, fainting, chills or fever, indigestion, loss of appetite, constipation, bruising, bleeding, swelling (oedema), weight loss, muscle aches, painful joints, back pain, decreased or loss of feeling or sensation such as numbness, especially in the skin, abnormal feeling or sensation such as tingling and 'pins and needles', sleeplessness, depression, confusion, nervousness, dry mouth, loss of memory, breathlessness, nosebleed, serious lung infection with fever, shortness of breath, cough, phlegm; hair loss, skin itching or discolouration, impotence, chest pain, reduction in blood platelets, which increases the risk of bleeding or bruising (thrombocytopenia), accumulation of fluid around the lungs or an increase in liver enzymes. Your doctor may do a blood test which may show an increase in your liver enzymes.

Rare side effects: may affect up to 1 in 1,000 people

Bleeding gums, weight gain, severe chest pain (angina pectoris), heart muscle disease, (signs include fatigue, chest pain and palpitations), enlarged heart, accumulation of fluid around the heart, painful spasm of the blood vessels on the heart (while resting, usually at night or early morning) (Prinzmetal angina), loss of coordination, difficulty in speaking, dry skin, migraine, visual disturbances or double vision, ringing in the ears, dizziness on standing up (especially when getting up from a sitting or lying position), increased need to pass water at night, pain, 'flu-like' symptoms, sleepiness, widening of blood vessels, inflammation of the large bowel (signs include: diarrhoea, usually with blood and mucus, stomach pain, fever), inflammation of the stomach (signs include: pain, nausea, vomiting), area of abnormal density in the lung, increased creatinine level in blood tests, which may be a sign of kidney problems.

The following side effects have been reported but it is not known exactly how often they occur:

- Potentially life-threatening, irregular heart beat (torsade de pointes);
- Inflammation of the liver, symptoms include nausea, vomiting, itching, yellowing of the skin and eyes, discoloration of stool and urine (hepatitis);
- Lung inflammation (signs include fever, coughing, difficulty breathing, wheezing; which causes scarring of the lungs) (allergic alveolitis, including interstitial lung disease, pneumonitis);
- Inflammation of the kidneys (tubulointerstitial nephritis).
- Stroke (see section 2).

Reporting of side effects

If you get any side effects talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme Website: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Anagrelide

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton, blister and bottle label after EXP. The expiry date refers to the last day of that month.

Blister Pack:

Do not store above 30°C.

HDPE Bottle pack:

Do not store above 30°C. Store in the original package in order to protect from moisture.

After first opening keep the bottle tightly closed and store at dry conditions.

If your doctor stops your medicine, do not keep any leftover capsules unless your doctor tells you to. Do not throw away any

medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information.

What Anagrelide contains

The active substance is anagrelide. Each capsule contains 0.5 mg anagrelide (as anagrelide hydrochloride monohydrate).

The other ingredients are:

Capsule contents: Lactose monohydrate (Pharmatose 200M), Cellulose, microcrystalline (Avicel PH 101), Croscarmellose sodium, Hydroxypropylcellulose (Klucel LF) & Magnesium stearate.

Capsule shell: Gelatin, Black iron oxide (E172) & Titanium dioxide (E171).

Printing ink: Shellac, Black Iron Oxide (E172) & Potassium hydroxide).

What Anagrelide looks like and contents of the pack

Size '4', hard gelatin capsule having grey opaque cap and white opaque body, imprinted '1453' on cap and '0.5 mg' on body with black ink containing white to off-white powder.

OPA/Al/PVC-Al blister pack:

Pack sizes: 84, 90 & 100 hard capsules

HDPE bottle pack: Available in white opaque round HDPE bottle pack with continuous thread closure. HDPE bottle pack contains silica gel as a desiccant.

Pack sizes: 100 hard capsules

Not all pack sizes may be marketed

Marketing Authorization Holder

Milpharm Limited
Ares Block, Odyssey Business Park
West End Road
Ruislip HA4 6QD
United Kingdom

Manufacturer

APL Swift Services (Malta) Ltd
HF26, Hal Far Industrial Estate, Hal Far,
Birzebbugia, BBG 3000
Malta

Or

Generis Farmacêutica, S.A.
Rua João de Deus, 19, Amadora
2700-487- Portugal

Or

Arrow Génériques
26 avenue Tony Garnier, Lyon – France,
69007-France

Or

Milpharm Limited
Ares Block, Odyssey Business Park, West
End Road,
Ruislip HA4 6QD
United Kingdom

This leaflet was last approved in 05/2022.