Possible side-effects

This medicine is unlikely to cause side-effects unless your child is allergic to the ingredients. If your child experiences any of the following, stop giving the medicine and seek immediate medical help:

 Unexplained wheezing or difficulty in breathing. swelling of the face, lips, tongue, neck or throat (severe allergic reactions).

If your child experiences any other side-effects or you are not sure about anything, talk to your doctor or pharmacist.

Reporting of side-effects:

If you get any side-effects, talk to your doctor, pharmacist or nurse. This includes any possible side-effects not listed on this carton. You can also report side-effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store.

By reporting side-effects you can help provide more information on the safety of this medicine.

Storing this medicine

Do not give this medicine after the expiry date shown on the bottle. Keep the bottle tightly closed. There are no special storage instructions for this medicine. Keep out of the sight and reach of children. Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment. Retain carton for full instructions.

Further information

What's in this medicine

The active ingredients in each 5 ml of medicine are: Glycerol 0.75 ml and Sucrose 1.7 g. Other Ingredients are: Citric acid monohydrate, sodium benzoate (E211), anthocyanin, blackcurrant flavour and juice (containing propylene glycol (E1520) and benzyl alcohol), liquid alucose and purified water.

What the medicine looks like

CALCOUGH® Children's Syrup is a clear, purple liquid available in 125 ml bottles.

+ Years



Children's Syrup

Glycerol & Sucrose



Blackcurrant Flavour



Children's Syrup

Glycerol & Sucrose

WHAT THIS MEDICINE IS FOR:

This medicine is a non-drowsy cough syrup, which relieves:

Dry coughs

ough:

Qo

ഗ

Õ

ă

Throat

Z

elie

- Tickly coughs
- Sore throats

BEFORE GIVING THIS MEDICINE TO YOUR CHILD: This product is suitable for most people, but a few people should not use it. If you are in any doubt contact your doctor or pharmacist. DO NOT GIVE YOUR CHILD THIS MEDICINE:

- If they are under 1 year old.
- If they are allergic to any of the ingredients. TALK TO YOUR DOCTOR OR PHARMACIST

If your child has an intolerance to some sugars. PREGNANCY AND BREAST-FEEDING You can take this medicine if you are

pregnant or breast-feeding. IF ANYONE HAS TAKEN TOO MUCH

If anyone has taken too much of this product, contact a doctor or your nearest Accident & Emergency Department (Casualty), taking this pack with you.

PL 15513/0392

Marketing Authorisation Holder: McNeil Products Limited, HP12 4EG, UK Manufacturer: McNeil Iberica S.L.U., 28805 Madrid, Spain.

This carton was prepared in October 2023. CALCOUGH® is a registered trade mark.

Rinse and recycle bottle, keep cap on. Recycle leaflet & carton.

UK Only

125 ml Θ

HOW TO USE THIS MEDICINE

Check the information below to see how much medicine to give.

- For oral use only. Do not give more medicine than the carton (and/or label) tells you to.
- Do not overfill the spoon. There is a double-ended spoon in the pack.
- Check the cap seal is not broken before first use. If it is, do not use this medicine.

Children under 1 vear

Do not give to children under 1 year old.

Children 1 year and over and adults

Age How much

Children 1-5 years Children 5 years and over and adults

One 5 ml spoonful Two 5 ml spoonfuls

(including the elderly)

Give the dose 3 or 4 times within 24 hours, if needed.

- · If your child does not feel better or starts to feel worse, tell your doctor or pharmacist.
- IF YOU FORGET TO GIVE THE MEDICINE

If you forget a dose, give the next dose when needed, DO NOT give a double dose.

SOME OF THE INGREDIENTS CAN CAUSE PROBLEMS

- This product contains 2.15 a alucose and 1.7 a sucrose in each 5 ml, which may be harmful to the teeth. If you have been told by a doctor that your child has an intolerance to some sugars, contact your doctor before using this medicine. This should be taken into account in patients with diabetes.
- Carbohydrate content of this medicine is 3.4 g per 5 ml.
- This medicine contains 3.53 mg propylene glycol (£1520) in each 5 ml dose, which is equivalent to 0.71 mg/ml.
- This medicine contains 10 ma sodium benzoate (E211) in each 5 ml which is equivalent to 2 g/ml.
- This medicine contains less than 1 mmol sodium (23 mg) per 5 ml, that is to say essentially 'sodium-free'.
- This medicine contains 0.000053 mg benzyl alcohol in each 5 ml which is equivalent to 0.000011 mg/ml. Benzyl alcohol may cause alleraic reactions. Ask your doctor or pharmacist for advice if you are pregnant or breast-feeding or if you have a liver or kidney disease. This is because large amounts of benzyl alcohol can build-up in your body and may cause side effects (called "metabolic acidosis"). Do not use for more than a week in young children (less than 3 years old), unless advised by your doctor or pharmacist. This is due to an increased risk due to accumulation in young children.

CONTENTS: Each 5 ml of oral liquid contains: Glycerol 0.75 ml, Sucrose **Calcough**) 1.7 g. Other ingredients include: Liquid glucose 2.15 q. E1520, E211, benzyl

Children's Syrup

Glycerol & Sucrose **Blackcurrant Flavour**

1+ Years

Effectively relieves dry, tickly coughs and sore throats

DOSAGE: For oral use only. Use the measuring spoon provided.

Children 1 to 5 years: One 5 ml spoonful. Children 5 years and over (including adults and the elderly): Two 5 ml

spoonfuls. Give the dose 3 or 4 times within 24 hours, if needed,

Not suitable for children under 1 year. KEEP OUT OF THE SIGHT AND REACH OF CHILDREN. 900230E

McNeil Products Limited. HP12 4EG, UK.

Manufacturer: McNeil Iberica S.L.U., 28805 Madrid, Spain.



PL 15513/0392

alcohol. Carbohydate

content 3.4 g per 5 ml.

Please tell your doctor or

pharmacist if your child

does not feel better or

Read and retain carton

for full instructions. Keep

the bottle tightly closed.

There are no special

storage instructions.

starts to feel worse.