

STADA

Verified

Max: 8pt

Max:

Date: 23.04.2021

Artwork Number: 1211304

Local Authority submission required: Yes

Covonia Dry Cough Sugar Free Formula

Mat.No: 92789652004_150ML_LBA

Dimensions: 73 x 90mm

Barcode: **5011309432619**

Datamatrix: 2789652004

Move to synthetic back label, safety updates for opioids and excipient guideline updated

3. Colour: Keyline (Non-Printing)

6. Colour: Varnish Area

1. Fonts Used: Helvetica

Point Size Used: Min:

Point Size Used: Min:

Point Size Used: Min:

Point Size Used: Min:

Point Size Used: Min: 7pt

4. Colour: Print Free Area (Non-Printing)
5. Colour: Absolute Print Free Area

Product Title:

Keyline Ref:

Pharmacode:

Reason for Change

1. Colour: Yellow

2. Colour: Black

7. Colour: 8. Colour:

9. Colour:

11.Colour:

2. Fonts Used:

3. Fonts Used:

4. Fonts Used:

5. Fonts Used:

Proof Number: 4

Operator: Artv

Approved by:



Peels from here

This product contains pholocodine monohydrate, which reduces the desire to cough. It relieves dry coughs which are without phlegm.

1. Before you use this medicine

Do not use the medicine if you or your child have....

• An allergy to any of the ingredients listed in section 3.

• Liver disease • Shallow breath or breathing difficulties • Chest problems or chest infections for a long time • An asthma attack.

• Taken MAOI's (monoamine oxidase inhibitors) for depression within the last two weeks.

• Ever developed a severe skin rash or skin that peels, blisters and/or sores in the mouth after taking this medicine or other related medicines.

• Or your child is under 12 years old.


■ Talk to your doctor if you or your child have....

• Kidney problems • Liver problems.


• A history of asthma or other breathing problems including bronchitis.

• A cough for a long time or a cough which produces lots of phlegm. • A history of drug abuse; pholocodine is an opioid and addiction is observed with opioids as a class. Serious skin reactions, including acute generalized exanthematous pustulosis (AGEP) have been reported with the use of this medicine. AGEP is a generalized, red, scaly rash with bumps under the skin and vesicles associated with fever. Most common location: mainly located on the folds of the skin, the trunk and the upper limbs. The highest risk of serious skin reactions occurrence is especially during the first week of treatment. If you develop a severe rash or any of these skin symptoms, stop taking this medicine and contact or see a doctor immediately.


Base ply



Top ply reverse



Middle ply reverse



Print layout diagram
90mm X 73mm 3Ply Multipeel
with Alternating Inset Corner Opening