Possible side-effects

This medicine is unlikely to cause side-effects unless your child is alleraic to the ingredients. If your child experiences any side-effects or you are not sure about anything, talk to your doctor or pharmacist.

Reporting of side-effects:

If you get any side-effects, talk to your doctor, pharmacist or nurse. This includes any possible side-effects not listed on this carton. You can also report side-effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/vellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store.

By reporting side-effects you can help provide more information on the safety of this medicine.

Storing this medicine

Keep out of the sight and reach of children. Do not store above 25°C. Keep the bottle tightly closed. Do not use after the end of the month shown as the expiry date on the packaging. Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment. Retain carton for full instructions.

Further information

What's in this medicine

The active ingredient is: Glycerol 0.75 ml Other ingredients are: Purified water, maltitol liquid (E965), citric acid, hydroxyethylcellulose, sodium citrate and sodium benzoate (E211). The flavouring is apple (containing propylene glycol (E1520)).

What the medicine looks like

CALCOUGH® Infant Syrup is a colourless liquid available in 125 ml bottles. It contains 0.75 ml of the active ingredient, glycerol, in every 5 ml of medicine.





Infant Syrup

Glycerol





Infant Syrup

Glycerol

WHAT THIS MEDICINE IS FOR:

CALCOUGH® Infant Syrup is a soothing syrup which relieves the symptoms of:

• Dry coughs

Ž

Qo

C

Relie

Tickly coughs

REFORE GIVING THIS MEDICINE TO YOUR CHILD:

This product is suitable for most people, but a few people should not use it. If you are in any doubt you should contact your doctor or pharmacist.

DO NOT GIVE YOUR CHILD THIS MEDICINE:

- If they are under 3 months old.
- If they are allergic to any of the ingredients.

PL 15513/0168

Marketing Authorisation Holder: McNeil Products Limited, HP12 4EG, UK. Manufacturer:

McNeil Iberica S.L.U., 28805 Madrid, Spain. This carton was revised in October 2023. CALCOUGH® is a registered trade mark.

UK Only



Rinse and recycle bottle, keep cap on. Recycle leaflet & carton.

125 ml C

HOW TO USE THIS MEDICINE

Check the information below to see how much medicine to give.

- For oral use only
- Do not give more medicine than the carton (and/or label) tells you to.

Babies under 3 months

Do not give to babies under 3 months.

Children from 3 months to 5 years Age How much

Children 3 months to 1 year One 5 ml spoonful Children 1 - 5 years Two 5 ml spoonfuls

- Give the dose 3 or 4 times within 24 hours. if needed
- If symptoms persist for more than 3 days. consult your doctor.

IF ANYONE HAS TAKEN TOO MUCH

If anyone has taken too much of this product, contact a doctor or your nearest Accident & Emergency Department (Casualty), taking this pack with you.

IF YOU FORGET TO GIVE THE MEDICINE If you forget a dose, give the next dose when needed. DO NOT give a double dose.

TALK TO YOUR DOCTOR OR PHARMACIST If your child has an intolerance to some sugars.

SOME OF THE INGREDIENTS CAN CAUSE PROBLEMS

- This product contains maltitol. If you have been told by your doctor that your child has an intolerance to some sugars, contact your doctor before using this product. This should be taken into account in patients with diabetes.
- · Maltitol may have a mild laxative effect. Calorific value 2.3 kcal/g maltitol.
- This medicine contains less than 1 mmol sodium (23 mg) per 5ml, that is to say essentially 'sodium-free'
- This medicine contains 10 ma sodium benzoate (E211) in each 5ml which is equivalent to 2ma/ml.
- This medicine contains 18.24 mg propylene glycol (E1520) in each 5ml which is equivalent to 3.65 mg/5ml.

Please consult your

more than 3 days.

doctor or pharmacist if symptoms persist for

Read and retain carton

for full instructions. Keep

bottle tightly closed. Do

not store above 25°C.

Calcough)

Infant Syrup

Glycerol

3+ Months

Soothes & relieves tickly coughs DOSAGE: For oral use only. Use the

measuring spoon provided. Children 3 months to 1 year: One 5 ml spoonful.

Children 1 to 5 years: Two 5 ml spoonfuls.

Give the dose 3 or 4 times within 24 hours, if needed, Not suitable for children under 3 months.

KEEP OUT OF THE SIGHT AND REACH OF CHILDREN. McNeil Products Limited, HP12 4EG, UK.

Manufacturer: McNeil Iberica S.L.U., PL 15513/0168 28805 Madrid, Spain.