



CORSODYL

Spray

chlorhexidine digluconate

Please read right through this leaflet before you start using this medicine.

This medicine is available without prescription but you still need to use Corsodyl Spray carefully to get the best results from it.

- Keep this leaflet, you may need to read it again.
- If you have any other questions, or if there is anything you do not understand, ask your dentist.

In this leaflet:

1. What Corsodyl Spray does
2. Check before you use Corsodyl Spray
3. How to use Corsodyl Spray
4. Possible side effects
5. How to store Corsodyl Spray
6. Further information

1. What Corsodyl Spray does

Corsodyl Spray is effective in the treatment of many common dental conditions. The active ingredient is chlorhexidine digluconate which is an antibacterial agent that reduces the formation of dental plaque and is used to:

- **Treat and prevent gum disease**
(gums may bleed when brushed)
- **Promote gum healing after dental surgery or treatment**
- **Control mouth ulcers**
- **Maintain mouth hygiene**
- **Manage oral thrush and denture sore mouth**

DEVCOMP-0005171



2. Check before you use Corsodyl Spray



Do not use:

- **If you have ever had an allergic reaction** to chlorhexidine digluconate or any of the other ingredients (listed in Section 6).
- **In children under 12 years** unless recommended by a dentist.
- **Anywhere other than in the mouth.**
Wash thoroughly if you get any in your eyes or ears.
- **Immediately after brushing your teeth.**
See Take special care with Corsodyl for more information.



Take special care with Corsodyl

- **Stop using the spray and seek immediate medical help** if you get a **rash, swelling of the lips, tongue, throat or face, soreness or irritation of the mouth or have difficulty breathing.**
See Section 4 for more information.
- Temporary **staining of the teeth and tongue** may occur.
See Section 4 for more information.
- You may notice a **change of taste or burning sensation.**
See Section 4 for more information.
- **Toothpastes can stop Corsodyl working.** Rinse your mouth thoroughly with water and wait 5 minutes after brushing your teeth before using Corsodyl. You can also use the spray at a different time of day.
- **Do not bleach fabrics** that have been in contact with Corsodyl.

Corsodyl Spray contains macrogolglycerol hydroxystearate

- Macrogolglycerol hydroxystearate may cause skin reactions.



Pregnancy and breast feeding

- You can use Corsodyl if you are **pregnant or breast feeding.**

3. How to use Corsodyl Spray



- **Contains alcohol.** If swallowed by a child seek **medical help.**
- **Do not exceed the stated dose.**
- Remove cap and fit the pump head to the bottle.
- **Use up to 12 pump actions twice daily** (each pump action delivers 0.14 ml of spray).



- **Spray tooth and gum surfaces, ulcers or other sore areas.** Remove pump head after use and replace cap.
- **Gum disease:** One month course (one to two spray packs) is recommended.
- **Mouth ulcers and oral thrush:** Continue to use for 2 days after healing has occurred.
- **Denture sore mouth:** Dentures should also be cleaned and then sprayed with Corsodyl Spray.
- **Dental surgery:** Use as directed by your dentist. If your symptoms persist, **see your dentist.**

4. Possible side effects

Corsodyl can have side effects, but not everybody gets them.

- **Stop using the spray and seek immediate medical help** if you get a **rash, swelling of the lips, tongue, throat or face** or **have difficulty breathing**. These effects may be symptoms of a severe allergic reaction which is very rare.
- **Stop using the spray and tell your dentist** if you get **irritation of the mouth, soreness or swelling of the inside of the cheeks**. These effects should go away when treatment is stopped.

Other side effects that may occur:

- **Temporary staining of the tongue.**
This disappears when treatment is stopped.
- **Temporary staining of the teeth.** This can normally be removed by brushing. You can avoid staining by not drinking tea, coffee or red wine – especially one hour after use, and by brushing daily with toothpaste (see Section 2). If staining continues, it can be removed with a scale and polish from your dentist or hygienist. You can clean your dentures with denture cleaner.
- You may notice a **change of taste or burning sensation** when you first use the spray. This usually goes away with continued use.
- If **peeling of the skin** inside the mouth occurs, stop using the product and the effect should go away.

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this

leaflet. You can also report side effects directly via the Yellow Card Scheme at www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Corsodyl Spray

Keep out of the sight and reach of children.

Do not use after the 'EXP' date shown on the pack. Store below 25°C. Protect from light.

6. Further information

Active ingredients: the spray contains Chlorhexidine Digluconate 0.2% w/v.

Other ingredients: Ethanol 7% v/v, Macrogolglycerol Hydroxystearate, Sorbitol, Peppermint Oil and Water.

Each bottle of Corsodyl Spray contains 60 ml of spray.

The marketing authorisation holder is GlaxoSmithKline Consumer Healthcare (UK) Trading Limited, Brentford, TW8 9GS, U.K. and all enquiries should be sent to this address.

The manufacturer is Farmaclair, 14200 Herouville Saint Clair, France.

Leaflet revised: August 2018.

Trade marks are owned by or licensed to the GSK group of companies.

DEVCOMP-0005171
L619004/04